


# CLENPIQ®

(sodium picosulfate, magnesium oxide, and anhydrous citric acid) Oral Solution

10 mg/3.5 g/12 g per 160 mL bottle


## HOW TO TAKE CLENPIQ: DRINK, HYDRATE, REPEAT SPLIT-DOSE REGIMEN

The Day Before  
10:00 AM



Drink 5 cups (40 oz) or more of clear liquids.\*  
Finish liquids over the next 5 hours.

The Day Before  
6:00 PM



Drink 4 cups (32 oz) or more of clear liquids.  
Finish liquids 2 hours before your colonoscopy  
or as advised by your doctor.



Hydration is important and it's part of the prep.  
Make sure to hydrate before, during, and after the prep.

### Prep Assistant

The time of your colonoscopy is \_\_\_\_\_


1

ON THE DAY BEFORE THE COLONOSCOPY, stop eating all solid food and dairy, and start hydrating by drinking clear liquids.



- Drink one bottle of CLENPIQ  
 Hydrate

Drink 5 or more cups (8 oz each) of clear liquid.



Start: 10:00 AM

Finish: \_\_\_\_\_

Check off the cups as you go!

2

- Drink the other bottle of CLENPIQ  
 Hydrate

Drink 4 or more cups (8 oz each) of clear liquid.



Start: 6:00 PM

Finish: \_\_\_\_\_

Check off the cups as you go!

3

\*After your first dose, if severe bloating, swelling, or stomach pain occurs, delay the second dose until the symptoms resolve. Please see reverse page for additional information about CLENPIQ, the importance of a clear liquid diet, and hydration.

#### INDICATION

CLENPIQ® is a prescription medicine used by adults and children 9 years and older to clean the colon before a colonoscopy. CLENPIQ cleans your colon by causing you to have diarrhea. Cleaning your colon helps your healthcare provider see the inside of your colon more clearly during your colonoscopy.

#### IMPORTANT SAFETY INFORMATION

- Do not take CLENPIQ if your healthcare provider has told you that you have serious kidney problems, a blockage in your intestine (bowel obstruction), an opening in the wall of your stomach or intestines (bowel perforation), a very dilated intestine (toxic megacolon), problems with the emptying of food and fluid from your stomach (gastric retention), or an allergy to any of the ingredients in CLENPIQ.
- CLENPIQ and other bowel preparations can cause serious side effects, including serious loss of body fluid (dehydration) and changes in blood salts (electrolytes) in your blood. These changes can cause abnormal heartbeats that may result in death, seizures (this can happen even if you have never had a seizure), or kidney problems. Your chance of having fluid loss and changes in blood salts with CLENPIQ is higher if you have heart problems, have kidney problems, or take water pills or non-steroidal anti-inflammatory drugs (NSAIDs).

See Important Safety Information continued on the back